

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

Genomics

journal homepage: www.elsevier.com/locate/ygeno

Methods

Improved variation calling via an iterative backbone remapping and local assembly method for bacterial genomes

Hongseok Tae^a, Robert E. Settlage^a, Shamira Shallom^a, Jasmin H. Bavarva^a, Dale Preston^b, Gregory N. Hawkins^b, L. Garry Adams^c, Harold R. Garner^{a,*}^a Virginia Bioinformatics Institute, Virginia Tech, Blacksburg, VA, USA^b Texas Animal Health Commission, State-Federal Diagnostic Laboratory, Austin, TX, USA^c Department of Veterinary Pathobiology, College of Veterinary Medicine, Texas A&M University, College Station, TX, USA

ARTICLE INFO

Article history:

Received 8 May 2012

Accepted 31 July 2012

Available online 10 August 2012

Keywords:

Brucella

Sequence assembly

Resequencing

Variant calling

Comparative genomics

Iterative mapping

ABSTRACT

Sequencing data analysis remains limiting and problematic, especially for low complexity repeat sequences and transposon elements due to inherent sequencing errors and short sequence read lengths. We have developed a program, ReviSeq, which uses a hybrid method composed of iterative remapping and local assembly upon a bacterial sequence backbone. Application of this method to six *Brucella suis* field isolates compared to the newly revised *B. suis* 1330 reference genome identified on average 13, 15, 19 and 9 more variants per sample than STAMPY/SAMtools, BWA/SAMtools, iCORN and BWA/PINDEL pipelines, and excluded on average 4, 2, 3 and 19 variants per sample, respectively. In total, using this iterative approach, we identified on average 87 variants including SNVs, short INDELs and long INDELs per strain when compared to the reference. Our program outperforms other methods especially for long INDEL calling. The program is available at <http://reviseq.sourceforge.net>.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

The genome sequences of microbial field isolates often contain a substantial number of loci different from the published references due to the high rate of mutation in bacterial replication (ca. 1/300 per genome per replication) [1]. Fortunately variant calling in bacterial genomes is relatively straightforward compared to that for eukaryotic studies because bacterial genomes are haploid. Incorrect variant calling in bacterial genomes is often caused by structural variants or incorrect mapping due to sequence variants in diverse repeat sequences including tandem repeats and transposon elements. Sequencing errors rarely cause incorrect variant calling because they are easily identified by designing the study to have a high depth of raw sequence coverage (i.e. $>20\times$). Variants occurring in repeat sequences can incorrectly fool mapping programs into assigning high quality scores to incorrectly mapped reads when the sequence reads from the repeat loci are significantly different from the reference sequence (e.g. length variation at two or more tandem repeat loci containing the same motif often causes incorrect mapping of sequence reads and high quality scores to the reads). This leads directly to invalid variant calls in repeat loci because the variation calling programs rely only on the mapping quality scores to filter out false positive variants from incorrectly mapped reads. Several programs have been developed to find structural variations such as insertions, deletions and copy

number variation, but they also have a limitation in searching for long (i.e. >8 bases) insertions or deletions when the number of incorrectly mapped sequences at a locus is high. An improved mapping post processing step is necessary to correct for this class of incorrect variant calls.

To address these issues in variant calling, we have developed ReviSeq which uses an iterative backbone remapping and local assembly method to generate and revise bacterial genome sequences from short sequence reads and a reference sequence. Previous iterative retrieval approaches used in several *de novo* assembly methods [2,3] are limited in application to resequencing analysis because they do not assemble contigs into large structural sequences, especially in or near low complexity repetitive sequences. iCORN, which uses an iterative mapping approach to revise a genome sequence, was developed for resequencing, but it does not correct long INDELs because iCORN's approach uses simple iterative mapping and does not benefit from local re-assembly [4]. Here, we report an advanced iterative remapping and local assembly approach which generates the revised whole genome sequence structure at each iteration based upon a backbone sequence structure.

We demonstrated the effectiveness of this approach for identifying accurate sequence variants found within the bacterial mutants of *Brucella* field isolates. *Brucella* is a gram-negative pathogenic bacteria that causes zoonotic disease in domestic animals [5] and has been designated as a category B priority pathogen. Consuming milk products from or having direct contact with infected animals may result in transmission to humans via penetration of skin or mucosal membranes [6]. At the start of resequencing analysis project, it is important to choose a suitable

* Corresponding author at: Virginia Bioinformatics Institute, Virginia Tech, Washington Street, MC0477, Blacksburg, VA 24061-0477, USA. Fax: +1 540 231 2606.

E-mail address: garner@vbi.vt.edu (H.R. Garner).

reference genome sequence against which high probability variants can be identified. The variation identified is a foundation for many downstream analyses. Due to the pathogenicity of *Brucella*, the results of variation detection are the basis for developing assays that are critical to the detection and mitigation of *Brucella*, as both a potential bioterrorism threat and as an infectious agent.

The *Brucella* genome is composed of two circular chromosomes of 2.1 Mbp and 1.2 Mbp. The first fully sequenced organism in the genus *Brucella* was *Brucella melitensis* biovar 1 which was published in 2002 [7]. Currently, the complete genome sequences of 11 additional *Brucella* organisms are publically available and many other genomes from other strains in the genus *Brucella* are in the process of being sequenced. Here we used the genome sequence of *Brucella suis* 1330 as a reference for detection of variants in six field isolates of *Brucella* collected from several different hosts that exhibited highly similar characteristics to *B. suis* 1330 in the 'gold standard' antibody diagnostic tests and biochemical tests [8]. Currently, two different versions of *B. suis* 1330 genome sequences are available. The original sequence was published in 2002 [9] and has been used as a reference in several resequencing studies [10,11], and the revised sequence of the 'same' sample was published recently [12].

The *Brucella* genome contains an 842 bp *IS711* transposon element [13] that is unique to *Brucella* and exists at several different locations in the genome. The published *B. suis* 1330 reference genomes have seven copies of *IS711*. If two or more close proximity variants exist within a transposon element, then this can lead to incorrect mapping of sequencing reads and therefore wrong variant calling in these regions.

The published *B. suis* 1330 genome sequences also have 10 loci containing 8-mer tandem repeats (≥ 3 motif copies) which are highly variable in its field isolate genomes. When the lengths of tandem repeats at these loci are dramatically different from the reference, the reads containing these elements can produce invalid alignments or be mapped to incorrect loci, leading again to incorrect variant calls.

2. Results

2.1. Overview of the variant calling results

Through this iterative remapping and local assembly approach, we identified an average of 87 variants in genomes of field isolate samples with respect to the reference sequence (Supplemental table 1). Conversely, traditional mapping approaches called approximately 70 variants from genome sequences of each field isolate with respect to the revised *B. suis* 1330 genome [12]. The largest differences observed between the reference genome and field isolates were observed within two long additional sequences (69 bp and 78 bp) in genomes of all isolates (Fig. 1), which were not detected by traditional mapping methods. Interestingly, since the sequences exist not only in genome sequences of all six field isolates but also in seven other sequenced *Brucella* species – *Brucella abortus* S19, *Brucella abortus* biovar 1, *Brucella canis* ATCC

23365, *Brucella melitensis* 16M, *Brucella microti* CCM 4915, *Brucella ovis* ATCC 25840 and *B. suis* ATCC 23445 – the sequences are likely to be deletions in the specific sample used to generate the *B. suis* 1330 reference sequences rather than new insertions in the field isolates or closely related species.

2.2. Comparison with other resequencing analysis pipelines

To address the limitation of the traditional resequencing methods in correcting mismapping/misalignment of sequence reads, variant calling results of ReviSeq were compared with that from the traditional resequencing analysis method which uses a simple mapping and variant calling pipeline. The most well known pipelines are combinations of BWA/SAMtools and STAMPY/SAMtools. Both mapping programs, BWA (ver. 0.5.9) [14] and STAMPY (ver. 1.0.13) [15], were executed with default options, and SAMtools [16] (mpileup) was used to generate pileup files. Since the purpose of this comparison was to evaluate the limitation of the methods in variant calling for haploid genomes due to incorrect read alignments, we used a simple filtering approach to call variants instead of using variant calling programs such as 'bctools' of SAMtools or SNVer [17]. The variants were called from the pileup files only if they were supported by at least 40% of reads covering each locus which were themselves covered by at least 10 reads. For an insertion, we applied $40\% \times (\text{read length} - \text{insertion length}) / \text{read length}$, considering the probability of a read completely covering the inserted sequence. Another program, iCORN (ver. 0.97), which uses an iterative mapping method, was also compared. To assess the reliability of variant calling of each pipeline, we generated a consensus sequence for the pipeline by replacing the reference bases with the variants identified by the pipeline using sample 13. Then we remapped sequence reads of sample 13 to the consensus sequence using BWA and counted the number of problematic reads including unmapped reads, reads with mismatches, clipped (partially aligned) reads, pair unmapped reads and long distance pairs (> 500 bases, an average distance was 290 bases with 25 as a standard deviation) in the mapping result. The ReviSeq pipeline shows the smallest percentages for all types of problematic reads, which suggests that it is the most reliable among all compared pipelines (Fig. 2).

Average numbers of variants called by BWA/SAMtools, STAMPY/SAMtools pipelines and iCORN were 78, 74 and 72, respectively (Table 1). Compared to the results of ReviSeq, STAMPY/SAMtools, BWA/SAMtools and iCORN predicted on average of 4, 2 and 3 more variants, and called on average of 13, 15 and 19 less variants per sample respectively. Each variant region inconsistent with the results of ReviSeq and its aligned reads were visually inspected. All three pipelines showed limitations in identifying long INDELs, and called several incorrect SNVs and short INDELs mainly in long INDELs loci (Fig. 3). We also tested the ABYSS [2] assembly/BWASW [18] mapping/SAMtools pipeline and the BWA/SNVer pipeline along with the local alignment program of GATK [19] applied to the

```

Deletion site on chromosome 1
Sample.no17_chr1 1331729 CATTGATCTGTGTCTCGGCGCGCGCATCGGCAAGGTCGATACCCGTTTCTCTCCGCGAC
B.suis_1330_chr1 1331615 CATTGATCTGTGTCTCGGCGCGCATCG-----

Sample.no17_chr1 1331787 TTCCCTCCGCGCAATTGGCATCATAATCGACGCGCATCGAAGATATCATTATCATCG
B.suis_1330_chr1 1331644 -----AAGATATCATTATCATCG

Deletion site on chromosome 2
Sample.no17_chr2 463067 CCGATGGTGGCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCTCTTGT
B.suis_1330_chr2 463046 CCGATGGTGGCGAGGTTGCCCTG-----

Sample.no17_chr2 463126 TTCAACCCGGCTGCGGATCACCTTTCGACATAAGCGCCCTGCTGTTTGTCTGCGGGCG
B.suis_1330_chr2 463069 -----CTGTTTGTCTGCGGGCG

```

Fig. 1. Two long deletion sites in the genome sequence of the *Brucella suis* 1330 'reference' sample. These loci were not detected by traditional mapping methods. The similarities of the highlighted sequences were possible sources of deletion events and may have contributed to misalignments.




Fig. 2. Percentages of problematic reads. Sequence reads from sample 13 were re-mapped by BWA to the each consensus sequence, in which all variants identified by the corresponding pipeline from sample 13 replaced the reference bases. The graphs show the percentages of problematic reads including unmapped reads, reads with mismatches, clipped reads, pair unmapped reads and long distance pairs (> 500 bases) in the mapping result for each consensus sequence. The ReviSeq pipeline shows the smallest percentage of the problematic reads in all comparisons.

BWA/SAMtools pipeline but did not observe significant improvement (Supplemental Table 2).

INDELs predicted by PINDEL [20], a program to search for structural variations, were also compared with our results. The mapping results of BWA were used as input data to PINDEL. To reduce false positive calls in the PINDEL results, INDELs supported by at least 10 reads were chosen for comparison. Compared to the results of the ReviSeq pipeline, PINDEL predicted an average of 19 more variants and called an average of 9 less variants per sample (Table 2). Many of insertions and deletions falsely called by PINDEL were predicted from mis-alignments around long INDELs. Interestingly, a few insertions and deletions attributable to possible chimeric DNA, mutants, reads from different strains or contaminants existing in the samples were predicted by PINDEL but not by our method. Since the loci were unique regions in the genome and the frequency of reads contributing to these INDEL predictions was much lower than that of the average read depth (lower than 20% of average depth), but consistent in all variant calling results, we concluded that they were not derived from mis-mapping or mis-alignment (mapped to correct position but partially misaligned) and were therefore appropriately not called by our method.

We additionally sequenced four 8-mer tandem repeat loci and two C homopolymer loci for samples 17, 22, 29, 34 and 35 using the Sanger method to confirm the variants called by our method and the other pipelines, which were different (Supplemental Table 3). Peaks in chromatograms for two C homopolymer loci at all samples were not clearly identified due to possible heterozygous alleles (Supplemental

Fig. 1). ReviSeq could correctly identify alleles of 9 loci among 16 loci containing variants (4 in sample 17, 3 in sample 22, 3 in sample 29, 3 in sample 34 and 3 in sample 35), while STAMPY/SAMtools, BWA/SAMtools and iCORN pipelines identified 1, 0 and 0 loci. ReviSeq could not correctly identify alleles from seven tandem repeat loci of which allele lengths were close to or longer than the read lengths (76 bases), and this limitation could be reduced with longer reads.

2.3. Comparison of variants among field isolates

All six field isolates have a similar number of variants when compared with the revised reference. Among them, the number of common variants in all isolates totaled 39, including 32 SNVs and 7 INDELs. The pairwise comparison of their sequence variants is illustrative of the number of common variants of each pair (Supplemental Table 4), which reveals the evolutionary relationship of the samples (Fig. 4). Specifically, samples 13 and 22 have an additional insertion site of an *IS711* element at position 1,578,904 of chromosome 1, which has not yet been reported in any of the previously sequenced *Brucella* species. The insertion site immediately follows the stop codon of a protein coding locus annotated as a ribose ABC transporter at the *B. suis* 1330 reference.

As samples 29 and 34 were isolated from bovine tissue derived from the same herd, variants in their genomes were almost identical except for one SNV and the lengths of three 8-mer tandem repeat loci. Interestingly, although samples 13 and 22 were isolated from different hosts (samples 13 from equine tissue and 22 from bovine milk),

Table 1

Comparison to other resequencing analysis pipelines in identifying variants (SNVs:INDELs).

Sample no.	ReviSeq	STAMPY/SAMTOOLS				BWA/SAMTOOLS				iCORN			
		com.	–	+	total	com.	–	+	total	com.	–	+	total
13	89 (60:29)	78 (57:21)	11 (3:8)	6 (1:5)	84 (58:26)	76 (57:19)	13 (3:10)	4 (1:3)	80 (58:22)	69 (57:12)	20 (3:17)	7 (5:2)	76 (62:14)
17	85 (64:21)	74 (62:12)	11 (2:9)	2 (1:1)	76 (63:13)	73 (62:11)	12 (2:10)	1 (1:0)	74 (63:11)	70 (60:10)	15 (4:11)	2 (1:1)	72 (62:10)
22	84 (57:27)	75 (56:19)	9 (1:8)	4 (3:1)	79 (59:20)	70 (56:14)	14 (1:13)	3 (3:0)	73 (59:14)	68 (56:12)	16 (1:15)	4 (4:0)	72 (60:12)
29	88 (63:25)	73 (60:13)	15 (3:12)	3 (2:1)	76 (62:14)	71 (60:11)	17 (3:14)	1 (1:0)	72 (61:11)	71 (60:11)	27 (3:14)	2 (1:1)	73 (61:12)
34	87 (62:25)	71 (59:12)	16 (3:13)	4 (2:2)	75 (61:14)	70 (59:11)	17 (3:14)	1 (1:0)	71 (60:11)	70 (59:11)	17 (3:14)	0 (0:0)	70 (59:11)
35	87 (64:23)	73 (62:11)	14 (2:12)	4 (2:2)	77 (64:13)	71 (62:9)	16 (2:14)	2 (1:1)	73 (63:10)	71 (62:9)	16 (2:14)	0 (0:0)	71 (62:9)

com.: commonly identified variants by ReviSeq and another pipeline.

–: (count of variants identified by ReviSeq) – (count of commonly identified variants).

+: (count of variants identified by another pipeline) – (count of commonly identified variants).

Sample.no17_chr2_463067

CCGATGGTGGCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

GTGCGCAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTT

CCGATGG GCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTG

CCGATGGTGG GCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGGAGG GATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCAGGTTG ATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGAGGTTGCC TAGCGGAAGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGAGGTTGCCCTGATGTTT AACGGGCAGAGGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCAGGTTGCCCTGATGTTTGTAGC AAGCGGCCTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCAGGTTGCCCTGATGTTTGTAGCGGAAGGG GCGTCTTGTTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCG TTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCT TTTTC AACCCGGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCGAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTG CCGCGCTGCGGATCACCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

CCGATGGTGGCAGGTTGCCCTGATGTTTGTAGCGGAAGGGCAGAGGCGGCCTCTTGTT GCGCTGCGGATCCTCCCTTTGCGACATAAGCGCCCTGCTGTTTTTGTCTGCGGGCG

$$\pm: (\text{count of variants identified by PINDEL}) - (\text{count of commonly identified variants}).$$




Fig. 4. Phylogenetic tree constructed from variation data illustrates possible evolutionary relationships of the samples. The distances in the tree were measured by counting the number of different variants between samples.

still remain unexplained, and accurate variant calls may help target appropriate mechanistic studies necessary to explain this aspect of the host-pathogen behavior of this species. To further enhance the utility of this approach in bacterial genome research, inversion and rearrangement of a genome sequence with respect to the backbone,

which are also common variations in bacterial genomes and remain challenging, will be addressed in future versions.

4. Materials and methods

4.1. Sample preparation and sequencing

The Texas Animal Health Commission (TAHC) provided *Brucella* samples from six field isolates that exhibited characteristics highly similar to *B. suis* 1330 in standard antibody and biochemical diagnostic tests [8] for carbon dioxide utilization, production of hydrogen sulfide and dye (thionin and basic fuchsin) sensitivity. The samples were collected from equine tissue (sample 13), porcine tissue (sample 17), bovine milk (sample 22), and bovine tissue (sample 29, 34 and 35). Genomic DNA for each sample was sequenced via the Illumina GAIIX sequencer and data acquired using the standard Illumina 101 (sample 13) and 76 (sample 17, 22, 29, 34, and 35) cycle paired-end protocols generating approximately 23,500,000 sequencing read pairs (47,000,000 reads) per sample.

4.2. Iterative remapping and local assembly to generate correct consensus sequences

The Iterative remapping and local assembly approach used by ReviSeq are illustrated in Fig. 5. This process was performed on each sample independently. To begin the analysis, we trimmed all low quality base calls from the ends of sequencing reads to remove base calls that have a high probability of being incorrect. Trimming at each read started from the end base of each read and stopped when




Fig. 5. Iterative remapping and local assembly approach. BWA was used for mapping, and ABYSS and CLCBio genomics workbench were used for *de novo* assembly. The iteration of mapping and local assembly continued until the two consensus sequences S_{i-1} and S_i converged.

a high quality base call ($\geq Q20$, Phred quality score) was encountered. If one of the two reads in a pair were shorter than 20 bases after removing the low quality base calls, both reads were excluded. We then mapped the reads to the revised reference genome, S_0 , using BWA. This approach resulted in more than 99.9% of the reads mapping to the reference, yielding average sequence coverage of $1460\times$ and $1048\times$ for the 101- and 76-cycle data, respectively.

A new consensus sequence, S_1 , was generated using SAMtools from the read mapping results. Concurrent with mapping, we also performed two independent *de novo* assemblies of the data using ABYSS (with k-mer 55) and the CLC genomics workbench (with default options) to enable detection of long insertion and deletion events, and possible genomic re-arrangements. After assembly, we aligned the resulting contig sequences and the new consensus sequence, S_1 , to the reference sequence, S_0 , using BWASW which is a mapping tool to align long sequences to a reference. A new consensus sequence, S_2 , was generated using the simple majority voting method from the mapping results of the *de novo* assembly contigs and S_1 . For this step, we gave higher priority to S_1 than contigs. As a result, if S_1 and contig sequences were different at a locus, S_1 was chosen unless two *de novo* assembly contig sequences were consistent. This step allows for detection of large INDEL variants that are not correctly detected by a simple mapping method.

We next mapped the sequencing reads to the new consensus sequence, S_2 , using BWA to generate another consensus sequence, S_3 . If consensus sequences S_2 and S_3 were different, the reads were mapped to S_3 to test whether all reads were correctly aligned to the same positions and a new consensus, S_4 , was generated from the newly aligned reads. The mapping/compared was iterated until S_{i-1} and S_i converged. The purpose of this iteration was to remap incorrectly mapped reads which would otherwise cause incorrect variant calling at *IS711* transposon elements.

Next, partially mapped reads (clipped reads) to S_i were counted to search for long INDELs which were not detected in prior alignment steps. When the number of partially mapped reads and abnormal pairs (distance of a pair $> \mu + 3\sigma$ or $< \mu - 3\sigma$, where μ is the mean and σ is the standard deviation of the distance between pairs) was more than 10% of total mapped reads at a position in S_i , the partially mapped reads were locally assembled by searching for exact matches only and their contig was subsequently aligned to S_i . If a long INDEL was detected, S_i was modified and used as a new reference sequence and resubmitted to the iterative remapping process until there was no change.

4.3. Variants calling from consensus sequences

The final consensus sequence of each sample was aligned to the reference sequence by BWASW. Due to the varying lengths of the 8-mer tandem repeat loci and other long INDELs such as a new insertion of an *IS711* element in the sample genome, BWASW could not map the whole query sequence to the reference sequence as a single alignment. Instead, it fragmented the query sequence into several pieces and aligned them to the reference separately. We extended the partial alignments adding INDEL information to merge them into one alignment and called the variants from the alignment. Further, the length variants (from 40 bases deletions to 80 bases insertions) of the 8-mer tandem repeat loci in field isolate genomes were fixed and confirmed by Sanger sequencing.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.ygeno.2012.07.015>.

Acknowledgments

This work was supported by the Medical Informatics and Systems Division director's funds; the U.S. Department of Homeland Security through the National Center of Excellence for Foreign Animal and Zoonotic Disease Defense at Texas A&M University [subaward 570636 from DHS 2007-ST-061-000002]; and the National Institute of Health, National Human Genome Research Institute, and 1000 Genomes Project Dataset Analysis grant [grant number 1U01HG-005719-01]. Genomic DNAs of *Brucella* field isolates were provided by TAHC (Texas Animal Health Commission) and Texas A&M University. The Core Laboratory Facility and the Data Analysis Core at the Virginia Bioinformatics Institute sequenced the genomic DNAs and assembled contigs, respectively.

References

- [1] J.W. Drake, B. Charlesworth, D. Charlesworth, J.F. Crow, Rates of spontaneous mutation, *Genetics* 148 (1998) 1667–1686.
- [2] J.T. Simpson, K. Wong, S.D. Jackman, J.E. Schein, S.J. Jones, I. Birol, ABYSS: a parallel assembler for short read sequence data, *Genome Res.* 19 (2009) 1117–1123.
- [3] I.J. Tsai, T.D. Otto, M. Berriman, Improving draft assemblies by iterative mapping and assembly of short reads to eliminate gaps, *Genome Biol.* 11 (2010) R41.
- [4] T.D. Otto, M. Sanders, M. Berriman, C. Newbold, Iterative Correction of Reference Nucleotides (iCORN) using second generation sequencing technology, *Bioinformatics* 26 (2010) 1704–1707.
- [5] M.K. Glynn, T.V. Lynn, Brucellosis, *J. Am. Vet. Med. Assoc.* 223 (2008) 900–908.
- [6] E.J. Young, An overview of human brucellosis, *Clin. Infect. Dis.* 21 (1995) 283–289.
- [7] V.G. DelVecchio, V. Kapatral, R.J. Redkar, G. Patra, C. Muijer, T. Los, N. Ivanova, I. Anderson, A. Bhattacharyya, A. Lykidis, G. Reznik, L. Jablonski, N. Larsen, M. D'Souza, A. Bernal, M. Mazur, E. Goltsman, E. Selkov, P.H. Elzer, S. Hagius, D. O'Callaghan, J.-J. Letesson, R. Haselkorn, N. Kyrpides, R. Overbeek, The genome sequence of the facultative intracellular pathogen *Brucella melitensis*, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 443–448.
- [8] G.G. Alton, L.M. Jones, R.D. Angus, J.M. Verger, Techniques for the Brucellosis Laboratory, INRA, Paris, 1988.
- [9] I.T. Paulsen, R. Seshadri, K.E. Nelson, J.A. Eisen, J.F. Heidelberg, T.D. Read, R.J. Dodson, L. Umayam, L.M. Brinkac, M.J. Beanan, S.C. Daugherty, R.T. Deboy, A.S. Durkin, J.F. Kolonay, R. Madupu, W.C. Nelson, B. Ayodeji, M. Kraul, J. Shetty, J. Malek, S.E.V. Aken, S. Riedmuller, H. Tettelin, S.R. Gill, O. White, S.L. Salzberg, D.L. Hoover, L.E. Lindler, S.M. Halling, S.M. Boyle, C.M. Fraser, The *Brucella suis* genome reveals fundamental similarities between animal and plant pathogens and symbionts, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 13148–13153.
- [10] J.-P. Lavigne, A.C. Vergunst, G. Bourg, D. O'Callaghan, The IncP island in the genome of *Brucella suis* 1330 was acquired by site-specific integration, *Infect. Immun.* 73 (2005) 7779–7783.
- [11] A.R. Wattam, K.P. Williams, E.E. Snyder, N.F. Almeida, M. Shukla, A.W. Dickerman, O.R. Crasta, R. Kenyon, J. Lu, J.M. Shalom, H. Yoo, T.A. Ficht, R.M. Tsolis, C. Munk, R. Tapia, C.S. Han, J.C. Detter, D. Bruce, T.S. Brettin, B.W. Sobral, S.M. Boyle, J.C. Setubal, Analysis of ten *Brucella* genomes reveals evidence for horizontal gene transfer despite a preferred intracellular lifestyle, *J. Bacteriol.* 191 (2009) 3569–3579.
- [12] H. Tae, S. Shalom, R. Settlege, D. Preston, L.G. Adams, H.R. Garner, Revised genome sequence of *Brucella suis* 1330, *J. Bacteriol.* 193 (2011) 6410.
- [13] S.M. Halling, F.M. Tatum, B.J. Bricker, Sequence and characterization of an insertion sequence, *IS711*, from *Brucella ovis*, *Gene* 133 (1993) 123–127.
- [14] H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform, *Bioinformatics* 25 (2009) 1754–1760.
- [15] G. Lunter, M. Goodson, Stampy: a statistical algorithm for sensitive and fast mapping of Illumina sequence reads, *Genome Res.* 21 (2011) 936–939.
- [16] H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, The sequence alignment/map format and SAMtools, *Bioinformatics* 25 (2009) 2078–2079.
- [17] Z. Wei, W. Wang, P. Hu, G.J. Lyon, H. Hakonarson, SNVer: a statistical tool for variant calling in analysis of pooled or individual next-generation sequencing data, *Nucleic Acids Res.* 39 (2011) e132.
- [18] H. Li, R. Durbin, Fast and accurate long-read alignment with Burrows–Wheeler transform, *Bioinformatics* 26 (2010) 589–595.
- [19] A. McKenna, M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernysky, K. Garimella, D. Altshuler, S. Gabriel, M. Daly, M.A. DePristo, The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data, *Genome Res.* 20 (2010) 1297–1303.
- [20] K. Ye, M.H. Schulz, Q. Long, R. Apweiler, Z. Ning, Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads, *Bioinformatics* 25 (2009) 2865–2871.